



# ASTRO-H

## **INSTRUMENT CALIBRATION REPORT RIGIDITY AND SAA ASTH-GEN-CALDB-SAA**

Version 0.1

Aug 12 2016

**ISAS/ GSFC**

**Prepared by:** Hitomi Science Data Center

## Table of Contents

1	Introduction .....	4
1.1	Purpose .....	4
1.2	Scientific Impact .....	4
2	Release CALDB 20160310 .....	4
2.1	Data Description .....	4
2.2	Data Analysis .....	4
2.3	Results .....	6

**CHANGE RECORD PAGE (1 of 1)**

DOCUMENT TITLE : Rigidity and SAA			
ISSUE	DATE	PAGES AFFECTED	DESCRIPTION
Version 0.1	March 2016	All	First Release
Version 0.2	Aug 2016	Sec 2.1 2.2	Add HXI high background regions

## Introduction

### 1.1 Purpose

This document has two different sections: one devoted to the rigidity (COR) parameter and one to the South Atlantic Anomaly (SAA).

### 1.2 Scientific Impact

COR and SAA are two parameters used in the screening process. Both information are calculated and written for each observation in the .ehk file. They are derived for values stored in CALDB.

## 2 Release CALDB 20160310

Filename	Valid data	Release data	CALDB Vrs	Comments
ah_gen_cor_20140101v001.fits	2014-01-01	20160310	001	First release
ah_gen_saa_20140101v002.fits	2014-01-01	20160812	004	Second release of this file

### 2.1 Data Description

1) The rigidity values in this file are calculated using the International Geomagnetic Reference Field (IGRF) model updated in 2015. This model may be used to derive map of COR for different year. The rigidity map used by Hitomi is calculated for the year 2016.0. This information is stored in the file “ah\_gen\_cor”.


2) The SAA information stored in CALDB determines the origin of the calculation of the satellite position with respect to the SAA, and contains the longitude and latitude of the vertices that define the SAA for each Hitomi instrument. The file contains 6 columns and 6 rows. The columns are associated with the different instruments (SXI SXS HXI1 HXI2 SGD1 SGD2) and the rows are the vertices that defined the SAA. The vertices are different for each instrument This information is stored in the 1<sup>st</sup> extension of the file “ah\_gen\_saa”.

3) After launch the HXI determined regions of high background that are used instead of the standard SAA (described in 2) to screen the data. These new regions are written in the 2<sup>nd</sup> extension of the file “ah\_gen\_saa” that contains the longitude and latitude of the vertices that define the HXI high background region. These new definitions are introduced with the Hitomi CALDB ver 4.

## 2.2 Data Analysis


1) Comparison between the rigidity maps derived from the IGRF 2015 model for different years and the rigidity maps used by ASCA and Suzaku are shown in Figure 1. It appears very clearly (upper right corner image) that Suzaku data (COR2, Year 2006) has a distortion at longitude around 0 and positive latitude not seen in the other maps. Figure 2 shows a projection along longitudinal variations for different fixed values of the latitude where the distortion is more clear ( upper panel green curve).

ASCA and Suzaku Rigidity Maps Compared with geomag.f Results  
 Longitude range -180 deg to +180 deg in steps of 15 deg  
 Latitude range -35 deg to +35 deg in steps of 5 deg  
 Altitude = 6948.142 km (570 km above equator), except for COR



**Figure 1: Grayscale renderings of rigidity maps computed with the code “geomag.f” for the epochs 2000.0, 2006.0, 2010.0, and 2015.0, as compared to the COR (ASCA- not dated) and COR2 (Suzaku) maps**

ASCA and Suzaku Rigidity Maps Compared with geomag.f Results



rhili, plot\_rigidity\_vs\_lon\_150701.ps, 1-Jul-2015 21:42:32.00

**Figure 2: Cuts along latitude +20 deg, 0 deg, and -20 deg, as a function of longitude, for the same set of rigidity maps**

2) Fig 3 show the SAA region used for the data screening. These regions are the same used on board and they are set the same for all instruments




Figure 3: SAA region defined prelaunch for all instruments

3) Fig 4 show the high background region defined for the HXI, these regions are used used for the data screening instead of the SAA . These regions include the SAA as described for all other instruments but also others that affect the detector and the data screening




Figure 4: HXI high background region

## 2.3 Results

1) The rigidity map in CALDB for Hitomi contains a file based on the data derived from the code, using the 2015 IGRF model and computed for year 2106.0

### 3 Previous Release CALDB 20160310

Filename	Valid data	Release data	CALDB Vrs	Comments
ah_gen_cor_20140101v001.fits	2014-01-01	20160310	001	First release
ah_gen_saa_20140101v001.fits	2014-01-01	20160310	001	First release