

Suzaku Observations of X-ray Brightest Fossil Group ESO3060170

Yuanyuan Su
University of Alabama



University of Alabama
Raymond White III
Jimmy Irwin



University of Michigan
Renato Dupke





MIT
Eric Miller



Tokyo University
Liyi Gu

Groups \neq Small scale clusters missing metals/baryons
Groups' shallow potential \rightarrow deep observations at large radii

Fossil groups: central dominant galaxy ≥ 2 R-mags
brighter than 2nd brightest galaxy w/in $0.5 r_{\text{vir}}$ (Jones et al. 2003)




Are fossil groups old, undisturbed systems?

Yes

- Simulations (Navarro 95)
X-ray observations (Khosroshahi 07) :
→ concentrated dark matter
- Optical isophotes &
X-ray atmospheres: undisturbed
last significant mergers long ago

No

- Lack of defined cool cores:
cores heated by recent
mergers (Sun 04)
- Ratio of SN Ia to SN II iron
mass fraction $99 \pm 1\%$ (outer)
 $85 \pm 2\%$ (center) (Dupke10)




Are fossil groups old, undisturbed systems?

Yes



- Simulations (Navarro 95)
X-ray observations (Khosroshahi 07) :
→ concentrated dark matter
- Optical isophotes &
X-ray atmospheres: undisturbed,
last significant mergers long ago

No

- Lack of defined cool cores:
cores heated by recent
mergers (Sun 04)





Suzaku observed brightest fossil group ESO 3060170 (PI: Y.Su)




$$L_X \sim 3.45 \times 10^{43} \text{ erg s}^{-1}$$


$$L_R \sim 3.28 \times 10^{11} L_{R\odot}$$








Iron mass to light ratio reaches cluster level




SN Ia/SN II ratio is consistent over 0-130 and 130-300 kpc

No central enhancement of SN Ia ejecta

Comparisons with other fossil groups



Comparisons with other fossil groups



At $0.3 R_{\text{vir}}$, ESO3060170 $f_b \sim 0.06$ (Sun+2004), RXJ 1559 $f_b \sim 0.1$ (Humphrey+2011)

Conclusions

- Hot core; Entropy is flat at large radii
- Iron mass to light ratio consistent with clusters
- Low optical luminosity at given L_x or M_{total}
→ deficiency in star/galaxy formation
- X-rays much more extended than optical
- Fossil group diversity: baryon fraction, abundance ratios, etc. (formation sequence)
→ Heterogeneous population

Ongoing work: XMM observations of fossil groups (PI: R.Dupke)